



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

SARS-CoV-2 seroprevalence among 7,950 health-care workers in the Region of Southern Denmark. (THEIJID-D-21-01420R1)

Sebastian von Huth , Søren Thue Lillevang , Birgit Thorup Røge ,
Jonna Skov Madsen , Christian Backer Mogensen ,
John Eugenio Coia , Sören Möller , Ulrik Stenz Justesen ,
Isik Somuncu Johansen

PII: S1201-9712(21)00731-1
DOI: <https://doi.org/10.1016/j.ijid.2021.09.023>
Reference: IJID 5706



To appear in: *International Journal of Infectious Diseases*

Received date: 3 May 2021
Revised date: 6 September 2021
Accepted date: 13 September 2021

Please cite this article as: Sebastian von Huth , Søren Thue Lillevang , Birgit Thorup Røge ,
Jonna Skov Madsen , Christian Backer Mogensen , John Eugenio Coia , Sören Möller ,
Ulrik Stenz Justesen , Isik Somuncu Johansen , SARS-CoV-2 seroprevalence among 7,950 health-
care workers in the Region of Southern Denmark. (THEIJID-D-21-01420R1), *International Journal of
Infectious Diseases* (2021), doi: <https://doi.org/10.1016/j.ijid.2021.09.023>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Highlights

- Health-care workers carry a pronounced risk of acquiring SARS-CoV-2 infection.
- The seroprevalence of SARS-CoV-2 antibodies was 2.1% among 7,950 participants.
- There is an occupational risk for infection by working on dedicated COVID-19 wards.

Journal Pre-proof

SARS-CoV-2 seroprevalence among 7,950 health-care workers in the Region of Southern Denmark. (THEIJID-D-21-01420R1)

Authors

Sebastian von Huth^{a,b,c,*}, Søren Thue Lillevang^d, Birgit Thorup Røge^c, Jonna Skov Madsen^{e,f}, Christian Backer Mogensen^{f,g}, John Eugenio Coia^{f,h}, Sören Möller^{i,j}, Ulrik Stenz Justesen^{i,k}, Isik Somuncu Johansen^{a,b,i}

Author affiliations

^aDepartment of Infectious Diseases, Odense University Hospital. J. B. Winsloews Vej 4, Indgang 18 Penthouse 2. sal, DK-5000 Odense C, Denmark

^bResearch Unit of Infectious Diseases, University of Southern Denmark, Denmark. J. B. Winsloews Vej 4, Indgang 18 Penthouse 2. sal, DK-5000 Odense C, Denmark.

^cDepartment of Internal Medicine, Lillebaelt Hospital, Kolding, Sygehusvej 24, DK-6000 Kolding, Denmark.

^dDepartment of Clinical Immunology, Odense University Hospital, J. B. Winsloews Vej 4, DK-5000 Odense C, Denmark.

^eDepartment of Biochemistry and Immunology, Lillebaelt Hospital, Vejle, Beriderbakken 4, DK-7100 Vejle, Denmark.

^fDepartment of Regional Health Research IRS, University of Southern Denmark, J. B. Winsloews Vej 19, 3, DK-5000 Odense C, Denmark.

^gDepartment of Emergency Medicine, Hospital Sønderjylland, Kresten Philipsens Vej 15, DK-6200 Aabenraa, Denmark.

^hDepartment of Clinical Microbiology, Hospital South West Jutland, Finsensgade 35, Bygning F, 1. sal, DK-6700 Esbjerg, Denmark.

ⁱDepartment of Clinical Research, University of Southern Denmark, J. B. Winsloews Vej 19, 3, DK-5000 Odense C, Denmark.

^jOPEN - Open Patient Data Explorative Network, Odense University Hospital, J. B. Winsløws Vej 4, DK-5000 Odense C, Denmark.

^kDepartment of Clinical Microbiology, Odense University Hospital, J.B. Winsloews Vej 21, 2, DK-5000 Odense C, Denmark.

*Corresponding author

Author email addresses

Sebastian von Huth: Sebastian.Lleicht.von.Huth@rsyd.dk

Søren Thue Lillevang: Soren.Lillevang@rsyd.dk

Birgit Thorup Røge: Birgit.Thorup.Roege@rsyd.dk

Jonna Skov Madsen: Jonna.Skov.Madsen@rsyd.dk

Christian Backer Mogensen: Christian.Backer.Mogensen@rsyd.dk

John Eugenio Coia: John.Eugenio.Coia@rsyd.dk

Sören Möller: Soren.Moller@rsyd.dk

Ulrik Stenz Justesen: Ulrik.Stenz.Justesen@rsyd.dk

Isik Somuncu Johansen: Isik.Somuncu.Johansen@rsyd.dk

Corresponding author

Sebastian von Huth, MD PhD

Email: Sebastian.Lleicht.von.Huth@rsyd.dk

Postal address: Department of Infectious Diseases, Odense University Hospital, J. B. Winsloews
Vej 4, Indgang 18 Penthouse 2. sal, DK-5000 Odense C, Denmark

Declarations of interest: none.

Word count (Abstract): 200

Word count (excluding Abstract, Tables, Acknowledgements and Reference list): 3189

Journal Pre-proof

Abstract

Objective

Health-care workers (HCWs) carry a pronounced risk of acquiring severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. We aimed to determine the seroprevalence and potential risk factors of SARS-CoV-2 infection among HCWs in the Region of Southern Denmark after the first pandemic wave in the spring of 2020.

Methods

Observational study conducted May-June 2020. SARS-CoV-2 IgG and IgM antibodies were measured in plasma. Participants were asked to complete a questionnaire consisting of demographic information, risk factors and COVID-19-related symptoms.

Results

A total of 7,950 participated. The seroprevalence of SARS-CoV-2 antibodies was 2.1% (95% CI 1.8-2.4). Seropositive participants were significantly older (mean age 48.9 years vs. 46.7 years in seronegative, $P=0.022$) and a higher percentage had experienced at least one symptom of COVID-19 ($P<0.001$). The seroprevalence was significantly higher among HCW working on dedicated COVID-19 wards (3.5%, OR 2.02 (95% CI 1.44-2.84). Seroprevalence was significantly related to 11-50 close physical contacts per day outside work (OR 1.54, 95% CI 1.07-2.22).

Conclusion

The prevalence of SARS-CoV-2 antibodies was low in HCWs. However, the occupational risk for contracting the infection is higher for those working on dedicated COVID-19 wards. Further, our

results imply that attention should be paid to occupational risk factors in planning pandemic preparedness.

Keywords

SARS-CoV-2; COVID-19; health-care workers; seroprevalence; antibodies; epidemiology.

Journal Pre-proof

Introduction

COVID-19 (Coronavirus disease 2019) has surged as an ongoing worldwide pandemic throughout 2020 (Park et al., 2020, Siordia, 2020). The first Danish cases were reported in late February 2020, and the initial spread of infections most likely originated from ski tourists returning from Northern Italy and Austria (Madsen et al., 2021). The first epidemic wave in Denmark peaked in late March and early April, with 9.2 patients admitted to hospital per 100,000 population (Madsen et al., 2021, Statens Serum Institut).

Several studies have demonstrated that health care workers (HCWs) have a significantly increased risk of contracting COVID-19, the infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (Galanis et al., 2021, Gomez-Ochoa et al., 2021). Current knowledge suggests that working as a healthcare professional poses an occupational risk of infection, and further raises concern about the occupational safety of HCWs. Previous studies have demonstrated a moderate-to-high risk of COVID-19 among HCWs throughout the world, including Italy (Felice et al., 2020), Spain (Suarez-Garcia et al., 2020), The Netherlands (Sikkema et al., 2020), Belgium (Schohy et al., 2021), Sweden (Rudberg et al., 2020), Norway (Molvik et al., 2021), Switzerland (Piccoli et al., 2021), France (Davidov et al., 2021), Brazil (Toniasso et al., 2021), USA (Barrett et al., 2020), and UK (Nguyen et al., 2020). Furthermore, HCWs are found to have a higher prevalence of antibodies against SARS-CoV-2 compared to the general population (Galanis et al., 2021, Gomez-Ochoa et al., 2021). Previous studies from Denmark found seroprevalences of SARS-CoV-2 among HCWs to be 3.4-4.04% (Iversen et al., 2020, Jespersen et al., 2020).

In general, the highest rates of seroprevalence among HCWs are found in countries and regions with widespread community infection and large numbers of COVID-19 patients admitted to the hospital (Barrett et al., 2020, Nguyen et al., 2020, Rudberg et al., 2020). Further, it has been demonstrated that disease severity and mortality among HCWs is generally lower than among

patients (Sahu et al., 2020). Compared to other professions, HCWs face a challenging task of wearing correct protective equipment while having close contact with COVID-19 patients and are often working in limited workspaces with close contact to colleagues (Agius et al., 2020).

Knowledge of the SARS-CoV-2 seroprevalence among HCWs is important in order to determine the occupational risk, and to understand and prevent spread of COVID-19 in health-care facilities, including hospitals. It is not known, to what extent HCWs acquire COVID-19 from contacts outside the hospital, and subsequently introduces COVID-19 at the wards, giving rise to in-hospital spread to both patients and between colleagues. Due to the nature of COVID-19, including the risk of serious illness and debilitating long-term sequelae, it is important to continue monitoring the prevalence among HCWs.

In the present study, we specifically collected information on general risk factors, which are known to pose a risk of contracting COVID-19 (Elmore et al., 2020). Further, we aimed to describe the prevalence of SARS-CoV-2 antibodies and potential risk markers for seroconversion among HCWs and administrative staff, with special focus on the travel history, extent of social contacts and other potential risk behaviours.

Methods

Study design

The Region of Southern Denmark covers approx. 12.000 km² and is inhabited by 1.2 million people (Region of Southern Denmark). The Region is administratively responsible for healthcare service, and runs 13 somatic hospitals and 12 psychiatric hospitals (Region of Southern Denmark). HCWs

and administrative staff in the Region of Southern Denmark (N=30,490 at the present time) were invited to participate on a voluntary basis. HCWs and administrative staff included both staff employed at somatic and psychiatric wards, prehospital and staff employed at general practitioners. The project was announced on the local intranets, and invitations to participate were sent online to the employee's state-provided personal and password-protected email system (e-Boks). Invitations were sent out in May 2020, and participants were allowed to accept participation until June 2020. All participants were asked to fill out a corresponding questionnaire, as described below. All employees were offered serological testing, regardless of their participation in the questionnaire. The project was registered with the Danish Data Protection Authorities (ref. no. 20/20627). The Regional Committees on Health Research Ethics for the Region of Southern Denmark evaluated the project and found that further registration and ethical permission was not necessary. The study was initiated by a group of senior scientists and supported by the national organization Danske Regioner ("Danish Regions").

Serological testing

Blood sampling was performed at the local hospital laboratory in designated EDTA blood collection tubes. SARS-CoV-2 IgG and IgM were measured in plasma with the lateral flow assay Livzon IgM/IgG LFT - Diagnostic Kit for IgM/ IgG Antibody to Corona Virus (Zhuhai Livzon Diagnostics, Inc., Zhuhai, China). The test is CE-IVD approved, and uses a colloidal gold immuno-chromatography technology to detect either IgM or IgG SARS-CoV-2 antibodies. The assay was performed according to the manufacturer's instructions as described by Nilsson et al. (Nilsson et al., 2021). The result of the assay was read by visual inspection by trained laboratory personnel 15 minutes after application of test material (one observer per test, but tests were performed by multiple laboratory technicians). Only tests in which the control line was visible were regarded as

valid. If the control line was not visible, the test was repeated. If no control line was visible when the test was repeated, the test was considered negative. If a line for IgM and/or IgG was observed, the test was defined as positive for that isotype of antibody.

Two batch numbers (CK2004150410 and CK2003100410) were used for the study. The batches were validated using a test panel of 600 blood donor samples from February 2018 and February 2019 (negative controls) and 150 samples from patient's previously tested PCR positive for SARS-CoV-2 (positive controls). The sensitivity of the two batches was 77.3% (CK2004150410) and 78% (CK2003100410), respectively, and the specificity was 99.3% (CK2004150410) and 98.7% (CK2003100410), respectively.

We used data on seroprevalence from healthy blood donors inhabiting the same geographical area as our study participants region (The Region of Southern Denmark), in order to compare results with the general population as described by Erikstrup et al. in (Erikstrup et al., 2021) and further personal communication with the authors.

Questionnaire

All participants were asked to fill out an online questionnaire in Danish on a secured platform. The questionnaire contained questions regarding employment data, demographics, information on chronic illness, travelling history, and symptoms of infection. The questionnaire was designed by the author group with inspiration from the questionnaire applied by Iversen et al. (Iversen et al., 2020), and was based on known risk factors for COVID-19 and other viral diseases. The questionnaire was proofread by both laymen and health-care workers not involved in the study. The complete questionnaire in Danish and translated to English is available in the Supplementary material.

Data handling and statistical analyses

Online questionnaires were archived on a secured online REDCap-based system provided by OPEN (Open Patient Data Explorative Network) (Harris et al., 2019, Harris et al., 2009). Serological data was merged with questionnaire data through the participants' social security number. Anonymized data was extracted by a dedicated data manager, who was not involved in the analysis and interpretation of the results.

The outcome investigated was seroprevalence reported as counts and proportions with exact binomial 95% confidence intervals (CI). The overall seroprevalence estimates were adjusted to mean test sensitivity and specificity of the two batches by the method suggested by Rogan and Gladen (Rogan and Gladen, 1978), and reported with Wald confidence intervals.

Associations with possible risk factors were investigated by univariate logistic regression reporting odds ratios (OR) with 95% CI and P-values for absence of association.

All analyses were performed in Stata 16.1. P-values below 0.05 were considered statistically significant.

Results

Study participant characteristics

In May 2020, all HCWs and administrative staff employed by the Region of Southern Denmark were invited to participate in the study. A total of 20,510 persons provided blood for serological testing. Of these, 7,950 (38.8%) provided questionnaire data. Only individuals with both serological and questionnaire data were included in the study.

The mean age of the participants was 46.7 years (SD 11.9 years, range 18-76 years) and 87.4% were female (Table I). Participants were from all professional groups, both with and without direct

patient contact, and included 37% nurses, 12% medical doctors, 13% administrative staff etc. (Table I). We did not find any significant differences between different professional groups regarding seropositivity.

Prevalence of SARS-CoV-2 antibodies

In total, 166 (2.1%, 95% CI 1.8%-2.4%) participants were found to have antibodies (either IgG, IgM or IgG and IgM) against SARS-CoV-2. Adjusting for sensitivity and specificity of the applied antibody test, this prevalence corresponds to an estimated true positive rate of 1.4% (95% CI 1.0-1.8%). Of the 166 positive samples, 143 (86.1%) were IgG positive, 111 (66.7%) were IgM positive and 88 (53.0%) were both IgG and IgM positive. There were no inconclusive test results.

The baseline characteristics stratified by antibody response are shown in Table I. The seropositive participants were significantly older (mean age 48.9 years vs. 46.7 years in seronegative). There was no significant difference according to sex ($P=0.477$) or body mass index between the two groups ($P=0.078$). Among the seropositive participants, a higher percentage had been PCR tested for SARS-CoV-2, compared to seronegative participants.

Self-reported symptoms

The association between self-reported symptoms and SARS-CoV-2 seroprevalence is shown in Table I and shown in detail including all reported symptoms in Supplementary Table I. Displaying at least one symptom of COVID-19 significantly increased the odds of having SARS-CoV-2 antibodies (OR 2.71, 95% CI 1.74-4.23).

Nearly all symptoms were significantly associated with an increased odds ratio for SARS-CoV-2 seropositivity, except nasal discharge or congestion, sore throat, conjunctivitis, and abdominal pain. We found that the most pronounced symptoms were loss of taste or smell (OR 15.22, 95% CI

10.73-21.58, $P<0.001$) and loss of appetite (OR 9.27, 95% CI 6.38-13.47, $P<0.001$). Further, fever (OR 6.02, 95% CI 4.40-8.23, $P<0.001$), chills (OR 5.34, 95% CI 3.72-7.68, $P<0.001$), chest pain (OR 4.59, 95% CI 2.88-7.31, $P<0.001$), coughing (OR 3.67, 95% CI 2.69-4.99, $P<0.001$) and shortness of breath (OR 3.47, 95% CI 2.29-5.27, $P<0.001$) were strongly associated with SARS-CoV-2 seropositivity. Reporting no symptoms was associated with a decreased OR of 0.33 (95% CI 0.21-0.53, $P<0.001$). 12.6% (20/163) of the participants with SARS-CoV-2 antibodies reported no symptoms prior to testing.

Reported type of work and occupational COVID-19 exposure

Table II describes the frequencies of positive antibody tests according to self-reported type of work, including work with direct patient contact. We found that working on dedicated COVID-19 wards was associated with a significant increased risk compared to those who did not work on dedicated wards (OR 2.02, 95% CI 1.44-2.84, $P<0.001$).

Number of close physical contacts, travel history and work from home

Self-reported potential work-related and personal risk factors for COVID-19 are shown in Table III. Seroprevalence was only significantly related to 11-50 close physical contacts outside work per day (OR 1.54, 95% CI 1.07-2.22, $P=0.021$), suggesting that crowding and multiple close contacts increase the risk of COVID-19. We did not find any association between working from home and a decreased prevalence of SARS-CoV-2 antibodies. Furthermore, our data does not support that travelling in the first months of 2020, prior to the first pandemic wave and corresponding lockdown, neither within nor outside Europe increased the risk of COVID-19 (Table III).

Geographical influence of SARS-CoV-2 seroprevalence

Geographical workplace and residence of participants are shown in Table III. We found that working in Southern Jutland significantly decreased the risk of SARS-CoV-2 seropositivity (OR 0.37, 95% CI 0.17-0.81, $P=0.013$), compared to working on the island of Funen (two hospitals; Odense University Hospital Odense and Svendborg). No other of the geographical workplaces included in our study were associated with increased or decreased risk.

The area of residence among participants did not seem to influence the risk of SARS-CoV-2 seropositivity.

The seroprevalence on different hospital sites is shown in Supplementary Table II. The seroprevalences on the larger hospitals (Odense University Hospital, Hospital of South West Jutland and Hospital of Lillebælt) were comparable, ranging from 2.1% to 2.8%. The SARS-CoV-2 seroprevalence on psychiatric departments was comparable to the larger somatic hospitals (2.4%). We found an overall low prevalence on Hospital Sønderjylland (0.4%), among general and specialist practitioners (0.0% and 0.9%, respectively) and prehospital staff (1.3%).

Alcohol and tobacco consumption, chronic comorbidities and SARS-CoV-2 seroprevalence

Association between SARS-CoV-2 seroprevalence and self-reported chronic diseases is depicted in Table IV. We did not find any association between chronic disease and SARS-CoV-2 seropositivity. Further, alcohol and tobacco consumption did not seem to be associated with SARS-CoV-2 seropositivity.

Discussion

The purpose of our study was to investigate the prevalence of SARS-CoV-2 antibodies in HCWs in the Region of Southern Denmark after the first pandemic wave of COVID-19 in the spring of 2020,

and to identify potential risk factors for infection. We found that among 7,950 participating HCWs, 2.1% (estimated true positive rate adjusted to test sensitivity and specificity 1.4%) were found to have SARS-CoV-2 antibodies. Our results suggest that working in dedicated COVID-19 wards poses an occupational risk of SARS-CoV-2 infection. In addition, the seropositive HCWs were significantly older. We also found that having a larger number of physically close contacts outside work increased the odds of seropositivity. Furthermore, our findings support previous studies regarding symptoms (Cascella et al., 2021, Hu et al., 2021), suggesting that displaying one or more symptom of COVID-19 increased the odds of seropositivity.

We find that participants that were PCR tested for SARS-CoV-2 had increased odds of seropositivity. Unfortunately, the result of the PCR test was not available, but in a setting with only limited access to PCR test for SARS-CoV-2, we presume that the participants had displayed symptoms of COVID-19.

The seroprevalence among HCWs and administrative staff in the Region of Southern Denmark in this paper is lower than those found in the Capital Region and the Central Denmark Region, where the seroprevalence was 4.04 and 3.4%, respectively (Iversen et al., 2020, Jespersen et al., 2020). This might reflect the distribution of the epidemic in Denmark, as the Region of Southern Denmark experienced one of the lowest overall prevalences of COVID-19. Furthermore, as the total number of infected persons and individuals admitted to the hospital in the other regions was larger than that of the Region of Southern Denmark, the risk of infection among HCWs was higher in the other regions.

As described previously, prevalence of SARS-CoV-2 antibodies among healthy blood donors in Denmark varied between different regions (Erikstrup et al., 2021), reflecting the overall regional variances in prevalence of COVID-19. In the Region of Southern Denmark during March and April

2020, the seroprevalence among blood donors was 1.74% (95% CI 0.43-2.16%), based on 4952 antibody tests. We found that the seroprevalence in HCWs and administrative staff was a little higher than in blood donors from the same period of time, which suggests that HCWs at the time were at increased risk of COVID-19, compared to the general population. This finding has been confirmed in other countries previously (Galanis et al., 2021, Gomez-Ochoa et al., 2021).

It is well established that HCWs are at greater risk of contracting SARS-CoV-2 infection compared to the general population (Grant et al., 2021, Rudberg et al., 2020), and that HCWs are prone to transmission of viral infection (Canova et al., 2020, McMichael et al., 2020, Ooi and Low, 2020, Oran and Topol, 2020, Ran et al., 2020, Riediker and Tsai, 2020, Sakurai et al., 2020, Wilson et al., 2020, Yu and Yang, 2020). Further, it has been demonstrated that HCWs are prone to infection despite vaccination (Bergwerk et al., 2021), and are able to transmit SARS-CoV-2 despite correct usage of personal protective equipment (Klompas et al., 2021).

Previous studies suggest that SARS-CoV-2 seropositivity is higher in HCWs performing patient-related work, in front-line HCWs, and HCWs working on dedicated COVID-19 wards (Grant et al., 2021, Iversen et al., 2020, Rudberg et al., 2020). Our findings are in line with this observation; however, we only see a significantly increased risk for HCWs working on dedicated COVID-19 wards, and no difference between HCWs with and without direct patient contact.

It is well established that asymptomatic carriers are able to spread infection (Ooi and Low, 2020, Rasmussen and Popescu, 2021). In an observational study from Canada, symptomatic SARS-CoV-2 infection in HCWs was found to be more common than asymptomatic, and only 0.50% of the asymptomatic participants were PCR-positive (Ferreira et al., 2021). In our study, 12.3% of seropositive HCWs did not report any symptoms, highlighting that although asymptomatic infections are infrequent, they may cause outbreaks among both patients and co-workers.

The main reported symptoms in our study were mild and included fever, nasal congestion, lethargy and headache. All reported symptoms were in line with previous publications and indistinguishable from other viral infections. This may partly explain the increased prevalence among HCWs compared to the general population; at the beginning of the epidemic, HCWs were less aware of the potential transmission of SARS-CoV-2 between colleagues, and perhaps attended work with mild symptoms. Furthermore, the PCR test capacity was limited during the first wave and mild symptoms were not an indication for a SARS-CoV-2 PCR test.

Previous studies have shown that healthcare assistants had higher prevalence of SARS-CoV-2 antibodies compared to other groups of HCWs, as these professional groups have most close-patient contact (Plebani et al., 2020, Rudberg et al., 2020). In our study, we did not find that a particular professional group had higher seroprevalence. This could suggest that viral transmission was not from patient to HCWs, but rather between HCWs. As personal protection equipment was not worn outside of separate patient rooms in Denmark during the first pandemic wave in the Spring of 2020, transmission between HCWs is very likely to have occurred.

There are some limitations to our study, which need to be taken into account. The sensitivity of the applied antibody test was rather low, increasing the risk of false negative results. However, we have adjusted the overall seroprevalence accordingly to the test sensitivity and specificity.

Our study setup allowed participants to have antibody tests to be performed without providing questionnaire data. This has led to the fact that a large number of participants did not provide questionnaire data. This could introduce a possible selection bias, which cannot be determined as

we have no information on the non-responders. Further, the questionnaire was designed so that questions could be left unanswered, which introduces some missing data.

Not all potentially important risk factors were considered in the questionnaire. Knowledge on risk behaviour both inside and outside health-care facilities, including compliance with personal protection equipment, hand sanitation habits and number of physical colleagues at work would have been of value.

Conclusion

In conclusion, we found low seroprevalence of SARS-CoV-2 among HCWs in the Region of Southern Denmark. However, working in dedicated COVID-19 wards posed a significantly higher occupational risk of SARS-CoV-2 infection. The majority of the seropositive HCWs had been symptomatic, which underlines the need for increased routine screening of HCWs in order to minimize the spread of the infection. Finally, increased attention should be paid to larger numbers of physically close contacts among HCWs during a pandemic.

Acknowledgements

The authors wish to thank all personnel involved in the serological analyses, including blood collection, handling, and analysis. Further, we wish to thank all study participants, and thank the hospital administrations and the Region of Southern Denmark for financial support.

Author contributions

Conceptualisation: Søren Thue Lillevang, Birgit Thorup Røge, Jonna Skov Madsen, Christian Backer Mogensen, John Eugenio Coia, Ulrik Stenz Justesen, Isik Somuncu Johansen

Statistical analysis: Sören Möller

Investigation: Sebastian von Huth, Ulrik Stenz Justesen, Isik Somuncu Johansen

Supervision: Ulrik Stenz Justesen, Isik Somuncu Johansen

Writing – original draft: Sebastian von Huth

Writing – review and editing: Sebastian von Huth, Sören Möller, Birgit Thorup Røge, Christian Backer Mogensen, John Eugenio Coia, Ulrik Stenz Justesen, Isik Somuncu Johansen.

All authors fulfil authorship criteria as defined by International Committee of Medical Journal Editors (ICMJE) and approve the submitted version of the article.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding source

This work was financially supported by Danish Regions (Danske Regioner) and the Region of Southern Denmark (Region Syddanmark). The funding sources were not involved in the study design, data collection, analysis, data interpretation, writing of the manuscript or the decision to publish.

Data availability

An anonymized dataset containing all data to obtain the results described in the present paper is available from the corresponding author on reasonable request.

References

- Agius RM, Robertson JFR, Kendrick D, Sewell HF, Stewart M, McKee M. Covid-19 in the workplace. *BMJ* 2020;370:m3577.
- Barrett ES, Horton DB, Roy J, Gennaro ML, Brooks A, Tischfield J, et al. Prevalence of SARS-CoV-2 infection in previously undiagnosed health care workers in New Jersey, at the onset of the U.S. COVID-19 pandemic. *BMC Infect Dis* 2020;20(1):853.
- Bergwerk M, Gonen T, Lustig Y, Amit S, Lipsitch M, Cohen C, et al. Covid-19 Breakthrough Infections in Vaccinated Health Care Workers. *N Engl J Med* 2021.
- Canova V, Lederer Schlapfer H, Piso RJ, Droll A, Fenner L, Hoffmann T, et al. Transmission risk of SARS-CoV-2 to healthcare workers -observational results of a primary care hospital contact tracing. *Swiss Med Wkly* 2020;150:w20257.
- Cascella M, Rajnik M, Aleem A, Dulebohn SC, Di Napoli R. Features, Evaluation, and Treatment of Coronavirus (COVID-19). *StatPearls*. Treasure Island (FL); 2021.
- David B, Gautier S, Riou I, Landowski S, Lawrence C, Thiebaut A, et al. The first wave of COVID-19 in hospital staff members of a tertiary care hospital in the greater Paris area: A surveillance and risk factors study. *Int J Infect Dis* 2021;105:172-9.
- Elmore R, Schmidt L, Lam J, Howard BE, Tandon A, Norman C, et al. Risk and Protective Factors in the COVID-19 Pandemic: A Rapid Evidence Map. *Front Public Health* 2020;8:582205.
- Erikstrup C, Hother CE, Pedersen OBV, Molbak K, Skov RL, Holm DK, et al. Estimation of SARS-CoV-2 Infection Fatality Rate by Real-time Antibody Screening of Blood Donors. *Clin Infect Dis* 2021;72(2):249-53.
- Felice C, Di Tanna GL, Zanusi G, Grossi U. Impact of COVID-19 Outbreak on Healthcare Workers in Italy: Results from a National E-Survey. *J Community Health* 2020;45(4):675-83.
- Ferreira VH, Chruscinski A, Kulasingam V, Pugh TJ, Dus T, Wouters B, et al. Prospective observational study and serosurvey of SARS-CoV-2 infection in asymptomatic healthcare workers at a Canadian tertiary care center. *PLoS One* 2021;16(2):e0247258.
- Galanis P, Vraika I, Fragkou D, Bilali A, Kaitelidou D. Seroprevalence of SARS-CoV-2 antibodies and associated factors in healthcare workers: a systematic review and meta-analysis. *J Hosp Infect* 2021;108:120-34.
- Gomez-Ochoa SA, Franco OH, Rojas LZ, Raguindin PF, Roa-Diaz ZM, Wyssmann BM, et al. COVID-19 in Health-Care Workers: A Living Systematic Review and Meta-Analysis of Prevalence, Risk Factors, Clinical Characteristics, and Outcomes. *Am J Epidemiol* 2021;190(1):161-75.

Grant JJ, Wilmore SMS, McCann NS, Donnelly O, Lai RWL, Kinsella MJ, et al. Seroprevalence of SARS-CoV-2 antibodies in healthcare workers at a London NHS Trust. *Infect Control Hosp Epidemiol* 2021;42(2):212-4.

Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.

Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42(2):377-81.

Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol* 2021;19(3):141-54.

Iversen K, Bundgaard H, Hasselbalch RB, Kristensen JH, Nielsen PB, Pries-Heje M, et al. Risk of COVID-19 in health-care workers in Denmark: an observational cohort study. *Lancet Infect Dis* 2020;20(12):1401-8.

Jespersen S, Mikkelsen S, Greve T, Kaspersen KA, Tolstrup M, Boldsen JK, et al. SARS-CoV-2 seroprevalence survey among 17,971 healthcare and administrative personnel at hospitals, pre-hospital services, and specialist practitioners in the Central Denmark Region. *Clin Infect Dis* 2020.

Klompas M, Baker MA, Griesbach D, Tucker R, Gallagher GR, Lang AS, et al. Transmission of SARS-CoV-2 from asymptomatic and presymptomatic individuals in healthcare settings despite medical masks and eye protection. *Clin Infect Dis* 2021.

Madsen LW, Lindvig SO, Rasmussen LD, Knudtzen FC, Laursen CB, Ovrehus A, et al. Low mortality of hospitalised patients with COVID-19 in a tertiary Danish hospital setting. *Int J Infect Dis* 2021;102:212-9.

McMichael TM, Currie DW, Clark S, Pogosjans S, Kay M, Schwartz NG, et al. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med* 2020;382(21):2005-11.

Molvik M, Danielsen AS, Grosland M, Telle KE, Kacelnik O, Eriksen-Volle HM. SARS-CoV-2 in health and care staff in Norway, 2020. *Tidsskr Nor Laegeforen* 2021;141(3).

Nguyen LH, Drew DA, Graham MS, Joshi AD, Guo CG, Ma W, et al. Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study. *Lancet Public Health* 2020;5(9):e475-e83.

Nilsson AC, Holm DK, Justesen US, Gorm-Jensen T, Andersen NS, Ovrehus A, et al. Comparison of six commercially available SARS-CoV-2 antibody assays-Choice of assay depends on intended use. *Int J Infect Dis* 2021;103:381-8.

- Ooi EE, Low JG. Asymptomatic SARS-CoV-2 infection. *Lancet Infect Dis* 2020;20(9):996-8.
- Oran DP, Topol EJ. Prevalence of Asymptomatic SARS-CoV-2 Infection : A Narrative Review. *Ann Intern Med* 2020;173(5):362-7.
- Park M, Cook AR, Lim JT, Sun Y, Dickens BL. A Systematic Review of COVID-19 Epidemiology Based on Current Evidence. *J Clin Med* 2020;9(4).
- Piccoli L, Ferrari P, Piumatti G, Jovic S, Rodriguez BF, Mele F, et al. Risk assessment and seroprevalence of SARS-CoV-2 infection in healthcare workers of COVID-19 and non-COVID-19 hospitals in Southern Switzerland. *The Lancet Regional Health - Europe* 2021;1:100013.
- Plebani M, Padoan A, Fedeli U, Schievano E, Vecchiato E, Lippi G, et al. SARS-CoV-2 serosurvey in health care workers of the Veneto Region. *Clin Chem Lab Med* 2020;58(12):2107-11.
- Ran L, Chen X, Wang Y, Wu W, Zhang L, Tan X. Risk Factors of Healthcare Workers With Coronavirus Disease 2019: A Retrospective Cohort Study in a Designated Hospital of Wuhan in China. *Clin Infect Dis* 2020;71(16):2218-21.
- Rasmussen AL, Popescu SV. SARS-CoV-2 transmission without symptoms. *Science* 2021;371(6535):1206-7.
- Region of Southern Denmark. Facts about the Region of Southern Denmark; Available from: <https://www.regionsyddanmark.dk/wm230808>. [Accessed March 19th 2021].
- Region of Southern Denmark. Sygehuse i Region Syddanmark; Available from: <https://www.regionsyddanmark.dk/wm205617>. [Accessed March 23rd 2021].
- Riediker M, Tsai DH. Estimation of Viral Aerosol Emissions From Simulated Individuals With Asymptomatic to Moderate Coronavirus Disease 2019. *JAMA Netw Open* 2020;3(7):e2013807.
- Rogan WJ, Gladen B. Estimating prevalence from the results of a screening test. *Am J Epidemiol* 1978;107(1):71-6.
- Rudberg AS, Havervall S, Manberg A, Jernbom Falk A, Aguilera K, Ng H, et al. SARS-CoV-2 exposure, symptoms and seroprevalence in healthcare workers in Sweden. *Nat Commun* 2020;11(1):5064.
- Sahu AK, Amrithanand VT, Mathew R, Aggarwal P, Nayer J, Bhoi S. COVID-19 in health care workers - A systematic review and meta-analysis. *Am J Emerg Med* 2020;38(9):1727-31.
- Sakurai A, Sasaki T, Kato S, Hayashi M, Tsuzuki SI, Ishihara T, et al. Natural History of Asymptomatic SARS-CoV-2 Infection. *N Engl J Med* 2020;383(9):885-6.

Scohy A, Gruson D, Simon A, Kabamba-Mukadi B, De Greef J, Belkhir L, et al. Seroprevalence of SARS-CoV-2 infection in health care workers of a teaching hospital in Belgium: self-reported occupational and household risk factors for seropositivity. *Diagn Microbiol Infect Dis* 2021;100(4):115414.

Sikkema RS, Pas SD, Nieuwenhuijse DF, O'Toole A, Verweij J, van der Linden A, et al. COVID-19 in health-care workers in three hospitals in the south of the Netherlands: a cross-sectional study. *Lancet Infect Dis* 2020;20(11):1273-80.

Siordia JA, Jr. Epidemiology and clinical features of COVID-19: A review of current literature. *J Clin Virol* 2020;127:104357.

Statens Serum Institut. COVID-19 i Danmark; Available from: <https://www.ssi.dk/covid19data>. [Accessed March 19th 2021].

Suarez-Garcia I, Martinez de Aramayona Lopez MJ, Saez Vicente A, Lobo Abascal P. SARS-CoV-2 infection among healthcare workers in a hospital in Madrid, Spain. *J Hosp Infect* 2020;106(2):357-63.

Toniasso SCC, Fernandes FS, Joveleviths D, Filho FFD, Takahasi AY, Baldin CP, et al. Reduction in COVID-19 prevalence in healthcare workers in a university hospital in southern Brazil after the start of vaccination. *Int J Infect Dis* 2021;109:283-5.

Wilson NM, Norton A, Young FP, Collins DW. Airborne transmission of severe acute respiratory syndrome coronavirus-2 to healthcare workers: a narrative review. *Anaesthesia* 2020;75(8):1086-95.

Yu X, Yang R. COVID-19 transmission through asymptomatic carriers is a challenge to containment. *Influenza Other Respir Viruses* 2020;14(4):474-5.

Table I: Baseline characteristics of the study participants

	Seronegative, N(%)	Seropositive, N(%)	OR (95% CI)	P-value
Total	7784 (97.8%)	166 (2.1%)		
Age (years) (mean, SD)	46.7 (11.9)	48.9 (11.9)	1.016 (1.002, 1.029)	0.022
Sex				
Female (N, %)	6803 (98.0%)	142 (2.0%)	1.00 (Ref.)	0.477
Male (N, %)	981 (97.6%)	24 (2.4%)	1.17 (0.76; 1.82)	
Body mass index (kg/m ²)				
<25	4137 (98.2%)	74 (1.8%)	1.00 (Ref.)	0.078
25-30	2290 (97.4%)	61 (2.6%)	1.49 (1.06; 2.10)	
≥30	1255 (97.9%)	27 (2.1%)	1.20 (0.77; 1.88)	
PCR tested for SARS-CoV-2	1511 (95.2%)	76 (4.8%)	3.61 (2.64; 4.94)	<0.001
≥ 1 symptom of COVID-19	5419 (97.4%)	143 (2.6%)	2.71 (1.74; 4.23)	<0.001
No symptoms	2271 (99.1%)	20 (0.9%)	0.33 (0.21; 0.53)	<0.001
Professional group				
Administrative staff	1058 (97.9%)	23 (2.1%)	1.00 (Ref.)	
Assisting nurse	435 (97.3%)	12 (2.7%)	1.27 (0.63; 2.57)	0.509
Laboratory technician	516 (99.0%)	5 (1.0%)	0.45 (0.17; 1.18)	0.104
Logistics staff	253 (96.6%)	9 (3.4%)	1.64 (0.75; 3.58)	0.218
Medical doctor	950 (97.4%)	25 (2.6%)	1.21 (0.68; 2.15)	0.513
Medical student	100 (100.0%)	0 (0.0%)	-	-
Nurse	2845 (97.6%)	69 (2.4%)	1.12 (0.69; 1.80)	0.653
Nursing student	107 (100.0%)	0 (0.0%)	-	-
Other staff with patient contact	740 (98.7%)	10 (1.3%)	0.62 (0.29; 1.31)	0.213
Other staff without patient contact	599 (98.2%)	11 (1.8%)	0.84 (0.41; 1.74)	0.649
Ph.D. student	31 (100.0%)	0 (0.0%)	-	-
Prehospital staff	70 (100.0%)	0 (0.0%)	-	-

Table II: Frequencies of positive antibody tests stratified according to type of work

	Seronegative, N(%)	Seropositive, N(%)	OR (95% CI)	P-value
Work without direct patient contact	1658 (97.8%)	37 (2.2%)	1.00 (Ref.)	0.750
Work involving direct patient contact	6044 (97.9%)	127 (2.1%)	0.94 (0.65; 1.36)	
Does not work on dedicated COVID-19 wards	3393 (96.7%)	116 (3.3%)	1.00 (Ref.)	<0.001
Work on dedicated COVID-19 wards	1309 (96.5%)	48 (3.5%)	2.02 (1.44; 2.84)	
Does not work in COVID-19 testing facilities	6393 (98.2%)	116 (1.8%)	1.00 (Ref.)	0.697
Work in COVID-19 testing facilities	284 (97.6%)	7 (2.4%)	1.16 (0.54; 2.51)	

Table III: Frequencies of positive antibody tests stratified according to exposure

	Seronegative, N(%)	Seropositive, N(%)	OR (95% CI)	P-value
Number of close physical contacts per day outside work				
<10	2515 (98.5%)	39 (1.5%)	1.00 (Ref.)	
11-50	4855 (97.7%)	116 (2.3%)	1.54 (1.07; 2.22)	0.021
51-100	309 (97.5%)	8 (2.5%)	1.67 (0.77; 3.61)	0.192
>100	36 (94.7%)	2 (5.3%)	3.58 (0.83; 15.40)	0.086
Working from home 4 weeks up to answered questionnaire				
No	6607 (97.8%)	146 (2.2%)	1.00 (Ref.)	
Yes, mainly	456 (98.9%)	5 (1.1%)	0.50 (0.20; 1.22)	0.125
Yes, partly	634 (98.1%)	12 (1.9%)	0.86 (0.47; 1.55)	0.609
Working from home in the period from mid-March to mid-April 2020				
No	6128 (97.8%)	135 (2.2%)	1.00 (Ref.)	
Yes, mainly	752 (98.7%)	10 (1.3%)	0.60 (0.32; 1.15)	0.126
Yes, partly	817 (97.8%)	18 (2.2%)	1.00 (0.61; 1.64)	1.000
Any travel history 2020				
No	5071 (98.0%)	102 (2.0%)	1.00 (Ref.)	
Yes	2713 (97.7%)	64 (2.3%)	1.17 (0.86; 1.61)	0.323
Any travel history 2020 (within Europe)				
No	5382 (98.0%)	110 (2.0%)	1.00 (Ref.)	
Yes	2402 (97.7%)	56 (2.3%)	1.14 (0.82; 1.58)	0.428
Travel history within 2020, but before serological testing				
Italy	138 (97.2%)	4 (2.8%)	1.37 (0.50; 3.74)	0.542
Austria	484 (98.2%)	9 (1.8%)	0.86 (0.44; 1.70)	0.674
Spain	308 (98.7%)	4 (1.3%)	0.60 (0.22; 1.63)	0.315
France	118 (99.2%)	1 (0.8%)	0.39 (0.05; 2.84)	0.355
Belgium	8 (80.0%)	2 (20.0%)	11.85 (2.50; 56.25)	0.002

Netherlands	42 (100.0%)	0 (0.0%)	-	-
Great Britain	88 (96.7%)	3 (3.3%)	1.61 (0.50; 5.14)	0.422
Other European countries	1575 (97.8%)	35 (2.2%)	1.05 (0.72; 1.54)	0.787
China	5 (100.0%)	0 (0.0%)	-	-
Iran	1 (100.0%)	0 (0.0%)	-	-
Asia (except China and Iran)	209 (98.6%)	3 (1.4%)	0.67 (0.21; 2.11)	0.490
Australia	20 (95.2%)	1 (4.8%)	2.35 (0.31; 17.63)	0.405
Africa	75 (97.4%)	2 (2.6%)	1.25 (0.31; 5.15)	0.754
North America	69 (100.0%)	0 (0.0%)	-	-
South America	27 (93.1%)	2 (6.9%)	3.50 (0.83; 14.86)	0.089
None of above	5004 (98.0%)	101 (2.0%)	0.86 (0.63; 1.18)	0.360
Geographic workplace				
Funen	3484 (97.9%)	74 (2.1%)	1.00 (Ref.)	
Lillebælt	2152 (97.6%)	53 (2.4%)	1.16 (0.81; 1.66)	0.416
Southern Jutland	888 (99.2%)	7 (0.8%)	0.37 (0.17; 0.81)	0.013
South West Jutland	918 (97.2%)	26 (2.8%)	1.33 (0.85; 2.10)	0.213
Other	342 (98.3%)	6 (1.7%)	0.83 (0.36; 1.91)	0.655
Area of residence				
Funen	3621 (97.8%)	83 (2.2%)	1.00 (Ref.)	
South Jutland	3746 (98.1%)	73 (1.9%)	0.85 (0.62; 1.17)	0.317
Region Central Jutland	281 (97.9%)	6 (2.1%)	0.93 (0.40; 2.15)	0.868
Region Northern Jutland	3 (100.0%)	0 (0.0%)	-	-
Region Zealand	18 (100.0%)	0 (0.0%)	-	-
Capital Region	24 (92.3%)	2 (7.7%)	3.64 (0.85; 15.64)	0.083
Outside Denmark	25 (96.2%)	1 (3.8%)	1.75 (0.23; 13.03)	0.587
Unknown	66 (98.5%)	1 (1.5%)	0.66 (0.09; 4.82)	0.683

Table IV: Frequencies of positive antibody tests stratified according to self-reported chronic diseases, smoking and alcohol consumption.

	Seronegative, N(%)	Seropositive, N(%)	OR (95% CI)	P- value
Self-reported chronic diseases				
Any chronic disease	1739 (97.9%)	37 (2.1%)	1.00 (0.69; 1.44)	0.987
Asthma	552 (97.9%)	12 (2.1%)	1.02 (0.56; 1.85)	0.946
Heart disease	117 (96.7%)	4 (3.3%)	1.62 (0.59; 4.44)	0.350
Hypertension	846 (98.0%)	17 (2.0%)	0.94 (0.56; 1.55)	0.797
COPD	61 (95.3%)	3 (4.7%)	2.33 (0.72; 7.50)	0.156
Kidney disease	27 (100.0%)	0 (0.0%)	-	-
Diabetes mellitus	139 (97.9%)	3 (2.1%)	1.01 (0.32; 3.21)	0.983
Immune deficiency	179 (96.8%)	6 (3.2%)	1.59 (0.70; 3.65)	0.270
Abdominal disease	156 (98.7%)	2 (1.3%)	0.60 (0.15; 2.43)	0.470
Alcohol consumption				
0 units/week	2665 (98.0%)	55 (2.0%)	1.00 (Ref.)	
1-7 units/week	4439 (97.9%)	95 (2.1%)	1.04 (0.74; 1.45)	0.832
8-14 units/week	507 (97.7%)	12 (2.3%)	1.15 (0.61; 2.16)	0.671
≥15 units/week	51 (100.0%)	0 (0.0%)	-	-
No answer provided	21 (100.0%)	0 (0.0%)	-	-
Tobacco use				
No, never	4755 (97.9%)	103 (2.1%)	1.00 (Ref.)	-
No, previous smoker	2174 (97.8%)	49 (2.2%)	1.04 (0.74; 1.47)	0.821
Yes, sometimes	319 (98.5%)	5 (1.5%)	0.72 (0.29; 1.79)	0.483
Yes, daily, <10 cigarettes/day	230 (98.3%)	4 (1.7%)	0.80 (0.29; 2.20)	0.669
Yes, daily, >10 cigarettes/day	184 (99.5%)	1 (0.5%)	0.25 (0.03; 1.81)	0.170
No answer provided	21 (100.0%)	0 (0.0%)	-	-